




Short Chain Fatty Acid Panel



Short-Chain Fatty Acids

Short-chain fatty acids (SCFAs) are produced in the colon by the gut microbiota. They are the end products of anaerobic fermentation of dietary fibers and protein/peptides in the small intestine. Their formation is the result of a complex interaction between diet, gut microbiota and host. SCFAs influence the physiology of the colon serving as energy sources by host cells and the intestinal microbiota as well as participating in different host-signaling mechanisms.

Applications

- ▶ Biopharmaceutical modulation of the host microbiome
- ▶ Bacterial product development
- ▶ Nutraceuticals and probiotics
- ▶ Dietary intervention
- ▶ Gut health and wellness
- ▶ Basic microbiome research

SCFA Panel	LLOQ*	
	Plasma/Serum	Feces
Acetic Acid (C2)	1000 ng/mL	50.0 µg/g
Propionic Acid (C3)	100 ng/mL	25.0 µg/g
Butyric Acid (C4)	25.0 ng/mL	25.0 µg/g
Isobutyric Acid (C4)	25.0 ng/mL	5.00 µg/g
Valeric Acid (C5)	25.0 ng/mL	5.00 µg/g
Isovaleric Acid (C5)	25.0 ng/mL	5.00 µg/g
2-Methyl butyric Acid (C5)	25.0 ng/mL	5.00 µg/g
Caproic Acid (C6)	50.0 ng/mL	1.00 µg/g
Lactic Acid	(upon request)**	

*Lower Limit of Quantitation (LLOQ) varies for each sample type.

** Available in select sample types only.

The panel is for non-GxP testing and is not for diagnostic use.

Analysis Method and Instrumentation

LC-MS/MS (Agilent 1290 UHPLC/Sciex QTrap 5500)

Sample Type and Required Amounts

Sample Type	Sample Requirement
Feces (wet weight)	200 - 500 mg
Plasma/Serum	100 - 150 µL
Others on request	

Contact us to get started
info@metabolon.com

+1 (919) 572-1711
metabolon.com
 617 Davis Drive, Suite 100, Morrisville, NC, 27560
 © 2021 Metabolon, Inc. All rights reserved. TA.SCFA.211111