




Salivary HETE

Oxylipins are signaling lipids that play key physiological roles including ovulation, initiation of labor, wound healing, kidney function, blood-vessel tone, blood coagulation, immune responses, and inflammation. They derive from the oxidation of polyunsaturated fatty acids (PUFA) through enzymatic or non-enzymatic free radical-mediated reactions, which can yield an array of lipid metabolites including eicosanoids, octadecanoids and docosanoids. Some of the most successful drug classes, such as the nonsteroidal anti-inflammatory drugs (NSAID), target the oxylipin modules.

Applications

- ▶ Gut health & wellness
- ▶ Immune system, infectious disease & COVID-19
- ▶ Inflammation & oxidative stress
- ▶ Cardiovascular diseases



Analyte	LLOQ
	Saliva
12-HETE	2.00 ng/mL

The panel is for non-GxP testing and is not for diagnostic use

Analysis Method and Instrumentation

LC-MS/MS (Agilent 1290 UHPLC/Sciex QTrap 5500)

Sample Type and Required Amounts

Sample Type	Sample Requirement
Saliva	250 - 300 μ L
Others on request	

Contact us to get started
info@metabolon.com

+1 (919) 572-1711

metabolon.com

617 Davis Drive, Suite 100, Morrisville, NC, 27560

© 2021 Metabolon, Inc. All rights reserved. TA.HETE.211111